

ASSIGNING THE COMPLICATED DISPERSED FLUORESCENCE SPECTRUM OF PhCCCN

JAMES H. THORPE, *Quantum Theory Project, University of Florida, Gainesville, FL, USA*; KHADIJA M. JAWAD, TIMOTHY S. ZWIER, *Department of Chemistry, Purdue University, West Lafayette, IN, USA*; JOHN F. STANTON, *Quantum Theory Project, University of Florida, Gainesville, FL, USA*.

The dispersed fluorescence spectrum from the S_1 origin of PhCCCN is reported. This molecule, which consists of a cyanoacetylene chain fused to an aromatic phenyl ring, is a potential candidate for astronomical detection. The high resolution spectrum was obtained under jet-cooled, gas-phase conditions with a rotational temperature of roughly 2 K, and appears to encode rich information about the vibronic coupling in this species. The experimentally observed peaks are assigned to emissions from the S_1 (\tilde{A}^1B_2) origin to a_1 vibrational levels of the ground state, which are dipole allowed and governed by FC factors, and to b_2 vibrational levels of the ground state, made possible by vibronic interactions between S_1 and S_2 (\tilde{B}^1A_1) states. The S_0 vibrational structure features a combined Fermi and Darling-Denison resonance near 950 cm^{-1} , which is accurately reproduced by diagonalization of an effective Hamiltonian containing the three a_1 states involved. A KDC Hamiltonian is employed to treat the vibronic b_2 features.